

## **Recombinant Insulin Like Growth Factor Binding Protein 7 (IGFBP7)**

Catalog No.: TP02585 50µg

**Sequence Information** 

Species: Human Gene ID:3490

Swiss Prot:Q16270 Synonyms:FSTL2; IGFBP-7v; MAC25; PSF;

TAF; IGFBP-rP1; MAC25 protein;

Prostacyclin-stimulating factor;

PGI2-stimulating factor; Tumor-derived

adhesion factor

Residues: Ser27~Leu282

SSSDTCGPCEPASCPPLPPLGCLLGETRDACGCCPMCARGEGEPCGGGGAGRGY

CAPGMECVKSRKRRKGKAGAAAGGPGVSGVCVCKSRYPVCGSDGTTYPSGCQLR

AASQRAESRGEKAITQVSKGTCEQGPSIVTPPKDIWNVTGAQVYLSCEVIGIPT

PVLIWNKVKRGHYGVQRTELLPGDRDNLAIQTRGGPEKHEVTGWVLVSPLSKED

AGEYECHASNSQGQASASAKITVVDALHEIPVKKGEGAEL

**Product Information** 

**Source:** Prokaryotic expression.

Host: E. coli

Tags: N-terminal His Tag

Subcellular Location: Secreted.

**Purity: >80%** 

**Traits:** Freeze-dried powder

**Buffer formulation:**PBS, pH7.4, containing 0.1% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 7.8

Predicted Molecular Mass: 30.2kDa

**Accurate Molecular Mass:** 35kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]



Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]




Figure 1. Gene Sequencing (Extract)






Figure 2. SDS-PAGE

## [ IMPORTANT NOTE ]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.