

## **Eukaryotic Cluster Of Differentiation 147 (CD147)**

Catalog No.: **TP05420** 50µg

**Sequence Information** 

Species: Human Gene ID:682

Swiss Prot: P35613 Synonyms: EMMPRIN; BSG; 5F7; M6; OK;

TCSF; Basigin(Ok Blood Group);

Collagenase stimulatory factor; OK

blood group antigen; Tumor cell-derived

collagenase stimulatory factor

Residues: Glu138~Ala323

EPGTVFTTVEDLGSKILLTCSLNDSATEVTGHRWLKGGVVLKEDALPGQKTEFK

VDSDDOWGEYSCVFLPEPMGTANIOLHGPPRVKAVKSSEHINEGETAMLVCKSE

SVPPVTDWAWYKITDSEDKALMNGSESRFFVSSSQGRSELHIENLNMEADPGQY

RCNGTSSKGSDQAIITLRVRSHLA

**Product Information** 

**Source:** Eukaryotic expression.

Host: 293F cell

Tags: N-terminal His Tag

Subcellular Location: Membrane.

**Purity: >90%** 

**Traits:** Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and

Proclin300.

Original Concentration: 200µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 4.8

Predicted Molecular Mass: 21.6kDa

Accurate Molecular Mass: 27-33kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[ STORAGE AND STABILITY ]



Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]




Figure 2. SDS-PAGE

## [ IMPORTANT NOTE ]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.