



Recombinant Activated Leukocyte Cell Adhesion Molecule (ALCAM)

Catalog No.: TP09062 50µg

Sequence Information

Species: Rat

Gene ID:79559

Swiss Prot:O35112

Synonyms:CD166; MEMD; Activated
Leucocyte Cell Adhesion
Molecule

Residues:Trp28~Lys527

WYTVNSAYGDTIVMPCRLDVPQNLMTGKWKYEKPDGSPVFIAPRSSTKKSQYDDVPEYKDRLSLSENYTSLINNAKIS
DEKRFVCMCLVTEDNVFEAPTLVKVFKQPSKPEIVNRAAFLETEQLKKLGDCISRDSYPDGNITWYRNGKVLQPDGEVS
ILFKKEIDPGTQLYTMSSLEYKTTKSDIQMPFTCSVTYYGPSGQKTIYSEQAIFDIYYPTQVTIQVLPKNAIKEGD
NITLQCLGNGNPPPEEFMFYLPGQAEGIRSSNTYTLTDVRRNATGDYKCSLIDQRNMAASTTITVHYLDLSLNPSGEVT
KQIGDTLPVSCTISASRNATVVMKDNIRLRSSPSFSSLHYQDAGNYVCETALQEVEGLKKRESLTLIVEGKPQIKMTK
KTDPSGLSKTIICHVEGFPKPAIQWTITGSGSVINQTEESPYINGRYYSKIIISPEENVTLTCTAENQLERTVNSLNVS
AISIPEHDEADDISDENREKVNDQAK

Product Information

Source: Prokaryotic expression.

Host: *E. coli*

Tags: N-terminal His Tag

Subcellular Location: Secreted.

Purity: >90%

Traits: Freeze-dried powder

Buffer formulation: PBS pH7.2, 0.01% sarcosyl, 5%Trehalose .

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 4.9

Predicted Molecular Mass: 57.3kDa

Accurate Molecular Mass: 58kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in PBS pH7.2 or ddH₂O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]




Figure 1. Gene Sequencing (Extract)




Figure 2 . SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.