

### Recombinant Human Aldolase C, Fructose Bisphosphate (ALDOC)

Catalog No.: TP01133 50µg

### **Sequence Information**

Species: Human Swiss Prot:P09972

### Gene ID:230

Synonyms:Fructose bisphosphate aldolase C , ALDC, ALDC, ALDO C, aldoc, ALDOC\_HUMAN, Aldolase 3, Aldolase C Fructose bisphosphateBrain type aldolase, Brain-type aldolaseFructoaldolase C.

#### Residues:Asp34~Ser281

DESVGSMAKRLSQIGVENTEENRRLYRQVLFSADDRVKKCIGGVIFFHETLYQK DDNGVPFVRTIQDKGIVVGIKVDKGVVPLAGTDGETTTQGLDGLSERCAQYKKD GADFAKWRCVLKISERTPSALAILENANVLARYASICQQNGIVPIVEPEILPDG DHDLKRCQYVTEKVLAAVYKALSDHHVYLEGTLLKPNMVTPGHACPIKYTPEEI AMATVTALRRTVPPAVPGVTFLSGGQSEEEAS **Product Information** 

Source: Recombinant expression. Host: *E.coli* Tags: N-terminal His-Tag Subcellular Location: Cytoplasm Purity: >90% Traits: Freeze-dried powder

**Buffer formulation:** PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.8

Predicted Molecular Mass: 30.8kDa

Accurate Molecular Mass: 31kDa as determined by SDS-PAGE reducing conditions.

# [ <u>USAGE</u> ]



Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

## [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [IDENTIFICATION]

KDa 180 130 70 55 40 35 25	
25 15 10	-

Figure 1. SDS-PAGE

# [<u>IMPORTANT NOTE</u>]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.