

Recombinant Human Cytoplasmic Protein NCK1 (NCK1)

Catalog No.: TP05091 100µg

Sequence Information

Species: Human Swiss Prot:P16333

Gene ID:4690

Synonyms:Cytoplasmic Protein NCK1; NCK adaptor Protein 1; Nck-1; SH2/SH3 Adaptor Protein NCK-Alpha; NCK1; NCK

Residues: Ala2~Ser377

AEEVVVVAKFDYVAQQEQELDIKKNERLWLLDDSKSWWRVRNSMNKTGFVPSNY

VERKNSARKASIVKNLKDTLGIGKVKRKPSVPDSASPADDSFVDPGERLYDLNM

PAYVKFNYMAEREDELSLIKGTKVIVMEKCSDGWWRGSYNGQVGWFPSNYVTEE

GDSPLGDHVGSLSEKLAAVVNNLNTGQVLHVVQALYPFSSSNDEELNFEKGDVM

DVIEKPENDPEWWKCRKINGMVGLVPKNYVTVMQNNPLTSGLEPSPPQCDYIRP

SLTGKFAGNPWYYGKVTRHQAEMALNERGHEGDFLIRDSESSPNDFSVSLKAQG

KNKHFKVQLKETVYCIGQRKFSTMEELVEHYKKAPIFTSEQGEKLYLVKHLS

Product Information

Source: Recombinant expression.

Host: E.coli

Tags: N-terminal His-Tag

Subcellular Location: Nucleus, Cytoplasm, Endoplasmic reticulum lumen

Purity: >90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.1

Predicted Molecular Mass: 46.4kDa

Accurate Molecular Mass: 46kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]



Reconstitute in ddH₂O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]




Figure 2. SDS-PAGE

[<u>IMPORTANT NOTE</u>]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.