

# Recombinant Acidic Nuclear Phosphoprotein 32 Family, Member A (ANP32A)

Catalog No.: **TP01171** 100µg

### **Sequence Information**

Species: Human Swiss Prot:P39687

### Gene ID:8125

Synonyms:Acidic leucine-rich nuclear phosphoprotein 32 family, member A; I1PP2A; LANP; MAPM; PHAP1; PHAPI; PP32; Mapmodulin.

#### Residues:Glu2~Asp175

EMGRRIHLELRNRTPSDVKELVLDNSRSNEGKLEGLTDEFEELEFLSTINVGLT

SIANLPKLNKLKKLELSDNRVSGGLEVLAEKCPNLTHLNLSGNKIKDLSTIEPL

KKLENLKSLDLFNCEVTNLNDYRENVFKLLPQLTYLDGYDRDDKEAPDSDAEGY

VEGLDDEEEDED

### **Product Information**

Source: Recombinant expression.

Host: E.coli

Tags: N-terminal His-Tag

Subcellular Location: Nucleus, Cytoplasm, Endoplasmic reticulum lumen,

Chromosome.

**Purity: >90%** 

Traits: Freeze-dried powder

**Buffer formulation:** PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

### Predicted isoelectric point: 4.5

Predicted Molecular Mass: 21.1kDa

Accurate Molecular Mass: 21kDa as determined by SDS-PAGE reducing conditions.

# [ <u>USAGE</u> ]

Reconstitute in  $ddH_2O$  to a concentration of 0.1-0.5 mg/mL. Do not vortex.



### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

# [ IDENTIFICATION ]




Figure 1. Gene Sequencing (Extract)

KDa	
180 130 100	=
70	
55	-
40	
35	
25	
15	
10	

### Figure 2. SDS-PAGE

## [ IMPORTANT NOTE ]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.