

Recombinant E3 ubiquitin-protein ligase TRIM38 (TRIM38)

Catalog No.: TP11129 100µg

Sequence Information

Species: Human

Swiss Prot: 000635

Gene ID:10475

Synonyms:TRI38

Residues:Met1-Asp465

MASTTSTKKMMEEATCSICLSLMTNPVSINCGHSYCHLCITDFFKNPSQKQLRQ ETFCCPQCRAPFHMDSLRPNKQLGSLIEALKETDQEMSCEEHGEQFHLFCEDEG QLICWRCERAPQHKGHTTALVEDVCQGYKEKLQKAVTKLKQLEDRCTEQKLSTA MRITKWKEKVQIQRQKIRSDFKNLQCFLHEEEKSYLWRLEKEEQQTLSRLRDYE AGLGLKSNELKSHILELEEKCQGSAQKLLQNVNDTLSRSWAVKLETSEAVSLEL HTMCNVSKLYFDVKKMLRSHQVSVTLDPDTAHHELILSEDRRQVTRGYTQENQD TSSRRFTAFPCVLGCEGFTSGRRYFEVDVGEGTGWDLGVCMENVQRGTGMKQEP QSGFWTLRLCKKKGYVALTSPPTSLHLHEQPLLVGIFLDYEAGVVSFYNGNTGC HIFTFPKASFSDTLRPYFQVYQYSPLFLPPPGD

Product Information

Source: Recombinant expression. Host: *E.coli* Tags: N-terminal His-Tag Subcellular Location: Secreted. Purity: >90% Traits: Freeze-dried powder Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300. Original Concentration: 200µg/mL Applications: Positive Control; Immunogen; SDS-PAGE; WB. (May be suitable for use in other assays to be determined by the end user.) Predicted isoelectric point: 7.1 Predicted Molecular Mass: 57.1kDa

Accurate Molecular Mass: 57kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in ddH_2O to a concentration of 0.1-0.5 mg/mL. Do not vortex.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]

KDa	
180 130 100	=
70	
55	
40	-
35 25	
25	_
15	
10	

Figure 1. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.