

Recombinant Gingipain R1 (rgpA)

Catalog No.: TP10358 100µg

Sequence Information

Species: PORGN

Swiss Prot: P28784

Gene ID:

Synonyms:CPG1

Residues:Tyr228-Ser720 YTPVEEKQNGRMIVIVAKKYEGDIKDFVDWKNQRGLRTEVKVAEDIASPVTANA IQQFVKQEYEKEGNDLTYVLLVGDHKDIPAKITPGIKSDQVYGQIVGNDHYNEV FIGRFSCESKEDLKTQIDRTIHYERNITTEDKWLGQALCIASAEGGPSADNGES DIQHENVIANLLTQYGYTKIIKCYDPGVTPKNIIDAFNGGISLVNYTGHGSETA WGTSHFGTTHVKQLTNSNQLPFIFDVACVNGDFLFSMPCFAEALMRAQKDGKPT GTVAIIASTINQSWASPMRGQDEMNEILCEKHPNNIKRTFGGVTMNGMFAMVEK YKKDGEKMLDTWTVFGDPSLLVRTLVPTKMQVTAPAQINLTDASVNVSCDYNGA IATISANGKMFGSAVVENGTATINLTGLTNESTLTLTVVGYNKETVIKTINTNG EPNPYQPVSNLTATTQGQKVTLKWDAPSTKTNATTNTARSVDGIRELVLLSVSD APELLRS

Product Information

Source: Recombinant expression. Host: *E.coli* Tags: N-terminal His and SUMO Tag Subcellular Location: Secreted. Purity: >90% Traits: Freeze-dried powder Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300. Original Concentration: 200µg/mL Applications: Positive Control; Immunogen; SDS-PAGE; WB. (May be suitable for use in other assays to be determined by the end user.) Predicted isoelectric point: 5.2 Predicted Molecular Mass: 69.9kDa Accurate Molecular Mass: 70kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]



Reconstitute in ddH₂O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]

KDa	
180 130 100	=
70	
55	-
40	-
35 25	
25	
15	-
10	

Figure 1. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.