

Recombinant Monoamine Oxidase B (MAOB)

Catalog No.: TP11338 100µg

Sequence Information

Species: Rat Swiss Prot:P19643 Gene ID: 25750

Synonyms:MAO-B; Amine oxidase

[flavin-containing] B

Residues:Gln206-Thr505

QERKFIGGSGQVSERIKDILGDRVKLERPVIHIDQTGENVVVKTLNHEIYEAKY

VISAIPPVLGMKIHHSPPLPILRNQLITRVPLGSVIKCMVYYKEPFWRKKDFCG

TMVIEGEEAPIAYTLDDTKPDGSCAAIMGFILAHKARKLVRLTKEERLRKLCEL

YAKVLNSQEALQPVHYEEKNWCEEQYSGGCYTAYFPPGILTQYGRVLRQPVGKI

FFAGTETASHWSGYMEGAVEAGERAAREILHAIGKIPEDEIWQPEPESVDVPAR

PITNTFLERHLPSVPGLLKLLGLTTILSAT

Product Information

Source: Recombinant expression.

Host: E.coli

Tags: N-terminal His-Tag

Subcellular Location: Mitochondrion outer membrane.

Purity: >90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 7.5

Predicted Molecular Mass: 37.1kDa

Accurate Molecular Mass: 38kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in ddH_2O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.



Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]

| KDa | |
|-------------------|---|
| 180 130 100 | |
| 70 | - |
| 55 | |
| 40 | |
| | - |
| 35 | |
| 25 | |
| | |
| | |
| | |
| 15 | |
| | |
| | |
| 10 | |
| 1U | and the second se |

Figure 1. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.