

Recombinant Poly ADP Ribose Glycohydrolase (PARG)

Catalog No.: TP10730 100μg

Sequence Information

Species: Human

Gene ID: 8505

Swiss Prot: Q86W56

Synonyms:

Residues: Ser448-Thr976

SPDKWLGTPIEEMRRMPPRCGIRLPLLPSANHTVTIRV DLLRAGEVPKPFPTH
YKDLWDNKHVVKMPCSEQNLYPVEDENGERTAGSRWEIQTALLNKFTRPQNLKD
AILKYNVAYSKKWDFTALIDFWDKVLEEAEAQHLYQSI LPDMVKIALCLPNICT
QPIPLLKQKMNHSITMSQE QIASLLANA FFTFPRR NAKM KSEYSSYPDINFNR
LFEGRSSRKPEKLKTLFCYFRRVTEKKPTGLVTFRQSLED FPEWERCEKPLTR
LHVTYEGTIEENGQGMLQVDFANRFVGGGVTSA GLVQEEIRFLINPELIISRLF
TEVLDHNECLIITGTEQYSE YTGYAETYRWSRS HEDG SERDDWQRRCTEIV AID
ALHFRRYLDQFVPEKMRRELNKAYCGFLRPGVSSENLSAVATGNW GCGA FGGDA
RLKALIQILAAAAAERDVYFTFGDSELMRDIYSMHIFLTERKLTVGDVYKLLL
RYYNEECRNCSTPGPDIKLYPFIYH AVE SCAETADHSGQRTGT

Product Information

Source: Recombinant expression.

Host: *E.coli*

Tags: N-terminal His Tag

Subcellular Location: Nucleus, Cytoplasm.

Purity: >90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.

Original Concentration: 200μg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 7.3

Predicted Molecular Mass: 64.8kDa

Accurate Molecular Mass: 61kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH₂O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]




Figure 1. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.